

# Solomon Practice Paper

## Core Mathematics 3E

Time allowed: 90 minutes

Centre: [www.CasperYC.club](http://www.CasperYC.club)

Name:

Teacher:

Question	Points	Score
1	5	
2	10	
3	11	
4	11	
5	12	
6	13	
7	13	
Total:	75	

How I can achieve better:

- 
- 
- 



Last updated: May 5, 2023




1. Express

$$\frac{2x^3 + x^2}{x^2 - 4} \times \frac{x - 2}{2x^2 - 5x - 3}$$

[5]

as a single fraction in its simplest form.



2. (a) Prove that, for  $\cos(x) \neq 0$ , [5]


$$\sin(2x) - \tan(x) \equiv \tan(x) \cos(2x).$$

(b) Hence, or otherwise, solve the equation [5]

$$\sin(2x) - \tan(x) = 2 \cos(2x).$$

for  $x$  in the interval  $0 \leq x \leq 180^\circ$ .

Total: 10



3.

$$f(x) = x^2 + 5x - 2 \sec(x), \quad x \in \mathbb{R}, \quad -\frac{\pi}{2} < x < \frac{\pi}{2}.$$

(a) Show that the equation  $f(x) = 0$  has a root in the interval  $[1, 1.5]$ . [2]

A more accurate estimate of this root is to be found using iterations of the form


$$x_{n+1} = \arccos(g(x_n)).$$

(b) Find a suitable form for  $g(x)$  and use this formula with  $x_0 = 1.25$  to find  $x_1, x_2, x_3$  and  $x_4$ . [6]  
Give the value of  $x_4$  to 3 decimal places.

The curve  $y = f(x)$  has a stationary point at  $P$ .

(c) Show that the  $x$ -coordinate of  $P$  is 1.0535 correct to 5 significant figures. [3]

Total: 11



4. (a) Differentiate each of the following with respect to  $x$  and simplify your answers. [6]

i.  $\sqrt{1 - \cos(x)}$


ii.  $x^3 \ln(x)$

(b) Given that [5]

$$x = \frac{y + 1}{3 - 2y},$$

find and simplify an expression for  $\frac{dy}{dx}$  in terms of  $y$ .

Total: 11




5. (a) Express  $\sqrt{3}\sin(\theta) + \cos(\theta)$  in the form  $R\sin(\theta + \alpha)$  where  $R > 0$  and  $0 < \alpha < \frac{\pi}{2}$ . [4]
- (b) State the maximum value of  $\sqrt{3}\sin(\theta) + \cos(\theta)$  and the smallest positive value of  $\theta$  for which this maximum value occurs. [3]
- (c) Solve the equation [5]

$$\sqrt{3}\sin(\theta) + \cos(\theta) + \sqrt{3} = 0,$$

for  $\theta$  in the interval  $-\pi \leq \theta \leq \pi$ , giving your answers in terms of  $\pi$ .

Total: 12



6. The function  $f$  is defined by

$$f(x) \equiv 3 - x^2, x \in \mathbb{R}, x \geq 0.$$


- (a) State the range of  $f$ . [1]
- (b) Sketch the graphs of  $y = f(x)$  and  $y = f^{-1}$  on the same diagram. [3]
- (c) Find an expression for  $f^{-1}$  and state its domain. [4]

The function  $g$  is defined by


$$g(x) \equiv \frac{8}{3 - x}, x \in \mathbb{R}, x \neq 3.$$

- (d) Evaluate  $fg(-3)$ . [2]
- (e) Solve the equation  $f^{-1}(x) = g(x)$ . [3]

Total: 13



7. Figure shows a graph of the temperature of a room,  $T^{\circ}\text{C}$ , at time  $t$  minutes.



The temperature is controlled by a thermostat such that when the temperature falls to  $12^{\circ}\text{C}$ , a heater is turned on until the temperature reaches  $18^{\circ}\text{C}$ . The room then cools until the temperature again falls to  $12^{\circ}\text{C}$ .

For  $t$  in the interval  $10 \leq t \leq 60$ ,  $T$  is given by

$$T = 5 + Ae^{-kt},$$

where  $A$  and  $k$  are constants.

Given that  $T = 18$  when  $t = 10$  and that  $T = 12$  when  $t = 60$ ,

(a) show that  $k = 0.0124$  to 3 significant figures and find the value of  $A$ , [6]

(b) find the rate at which the temperature of the room is decreasing when  $t = 20$ . [4]

The temperature again reaches  $18^{\circ}\text{C}$  when  $t = 70$  and the graph for  $70 \leq t \leq 120$  is a translation of the graph for  $10 \leq t \leq 60$ .

(c) Find the value of the constant  $B$  such that for  $70 \leq t \leq 120$  [3]

$$T = 5 + Be^{-kt}.$$

Total: 13

