


You will need to use a calculator for this worksheet

1



The diagram shows the curve $y = x^2$ which passes through the point $A(1, 1)$ and the point B .

- a Copy and complete the table to find the gradient of the chord AB when the x -coordinate of B takes each of the given values.

x -coordinate of B	y -coordinate of B	gradient of AB
2	4	$\frac{4-1}{2-1} = 3$
1.1	1.21	
1.01		
1.001		

- b Suggest a value for the gradient of the tangent to the curve $y = x^2$ at the point $(1, 1)$.
 c Repeat part a using 0, 0.9, 0.99 and 0.999 as the x -coordinates of B and comment on your answer to part b.

2 Use a similar table of values to that in question 1 to find a value for the gradient of the tangent to the curve $y = x^2$ at the point A when A has the coordinates

- a $(2, 4)$ b $(4, 16)$ c $(1.5, 2.25)$ d $(-3, 9)$

3 a Using your answers to questions 1 and 2, suggest an expression in terms of x for the gradient of the curve $y = x^2$ at the point (x, y) .

- b Write down the gradient of the curve $y = x^2$ at the points

- i $(6, 36)$ ii $(2.4, 5.76)$ iii $(-3.2, 10.24)$

4 By considering the gradient of a suitable sequence of chords, find a value for the gradient of each curve at the given point.

- a $y = x^4$ at $(1, 1)$ b $y = x^2 - 5x + 3$ at $(2, -3)$
 c $y = \sqrt{x}$ at $(4, 2)$ d $y = \frac{2}{x}$ at $(2, 1)$


5 a By considering the gradient of a suitable sequence of chords, find a value for the gradient of the curve $y = x^3$ at the points

- i $(1, 1)$ ii $(2, 8)$ iii $(3, 27)$

- b Suggest an expression of the form kx^n for the gradient of the curve $y = x^3$ at the point (x, y) .

c Find the gradient of the curve $y = x^3$ at the points

- i $(4, 64)$ ii $(-2, -8)$ iii $(1.5, 3.375)$



1 Differentiate with respect to x

a x^2 **b** x^4 **c** x **d** x^9 **e** x^{-3} **f** x^{-1}
g $4x^2$ **h** $7x$ **i** $2x^5$ **j** 3 **k** $8x^{-2}$ **l** $11x^{-4}$

2 Find $\frac{dy}{dx}$

a $y = x^5 + x^2$ **b** $y = x + x^3$ **c** $y = x^4 + 2$ **d** $y = x^6 - 2x$
e $y = 6x^3 + 5x^{-2}$ **f** $y = x^2 - 4x + 1$ **g** $y = x^{-1} - x^{-5}$ **h** $y = 4x^3 + 3x^{-4}$

3 Differentiate with respect to t

a t^6 **b** $5t^{-3}$ **c** $t^{\frac{1}{2}}$ **d** $t^{\frac{2}{3}}$ **e** $\frac{3}{4}t^2$ **f** $8t^{\frac{1}{4}}$
g $2t^{\frac{7}{2}}$ **h** $t^{-\frac{1}{5}}$ **i** $\frac{1}{2}t^{\frac{6}{5}}$ **j** $t^{-\frac{3}{2}}$ **k** $12t^{-\frac{5}{4}}$ **l** $\frac{1}{6}t^{\frac{4}{3}}$

4 Find $f'(x)$

a $f(x) = 2x + \frac{1}{3}x^6$ **b** $f(x) = x^{\frac{3}{2}} - 5$ **c** $f(x) = x + 4x^{\frac{1}{2}}$ **d** $f(x) = 6x^{\frac{5}{3}} - x^{-4}$
e $f(x) = 7 + x^{-\frac{4}{5}}$ **f** $f(x) = 2x^{\frac{1}{6}} + x^{\frac{3}{4}}$ **g** $f(x) = 3x^{-1} - 5x^{-\frac{3}{2}}$ **h** $f(x) = 2 - 7x^{-1} + x^{-\frac{8}{3}}$

5 Find $\frac{dy}{dx}$

a $y = \sqrt{x}$ **b** $y = 4 - \frac{1}{x}$ **c** $y = 3x^2 + \sqrt[3]{x}$ **d** $y = 9x + \frac{3}{x}$
e $y = \frac{1}{4x} - \frac{1}{x^2}$ **f** $y = \frac{6}{\sqrt[4]{x}}$ **g** $y = \sqrt{x^5}$ **h** $y = 8\sqrt{x} + \frac{4}{3x^2}$

6 Find $\frac{ds}{dt}$


a $s = t(t + 3)$ **b** $s = (t - 2)^2$ **c** $s = 5t(t^3 + 4t)$ **d** $s = t^2(7t - t^{-1})$
e $s = (t + 1)(t + 6)$ **f** $s = (t - 4)(t + 2)$ **g** $s = t(t^4 + 3t^2 + 9)$ **h** $s = t(t - 1)(2t - 3)$

7 Find $\frac{dy}{dx}$


a $y = \sqrt{x}(x - 4)$ **b** $y = \frac{x^3 - 2x}{x}$ **c** $y = \frac{4x^3 + x}{x^2}$ **d** $y = \frac{x + 3}{\sqrt{x}}$
e $y = \frac{4 - x^3}{2x}$ **f** $y = \frac{5 + \sqrt{x}}{x^2}$ **g** $y = \frac{9x - 2}{3x}$ **h** $y = \frac{8x + x^3}{4\sqrt{x}}$

8 In each case, find $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$.

a $y = 4x^2 - x + 3$ **b** $y = x^3 + 5x^2 + 2x - 6$ **c** $y = 8 - \frac{2}{x}$
d $y = 2x^4 + 3x^2 - 9$ **e** $y = \frac{3x^6 - 4}{x^2}$ **f** $y = 6x^{\frac{1}{2}} - x^{-\frac{1}{2}}$



- 1 Find the gradient at the point with x -coordinate 3 on each of the following curves.
- a** $y = x^3$ **b** $y = 4x - x^2$ **c** $y = 2x^2 - 8x + 3$ **d** $y = \frac{3}{x} + 2$
- 2 Find the gradient of each curve at the given point.
- a** $y = 3x^2 + x - 5$ (1, -1) **b** $y = x^4 + 2x^3$ (-2, 0)
- c** $y = x(2x - 3)$ (2, 2) **d** $y = x^2 - 2x^{-1}$ (2, 3)
- e** $y = x^2 + 6x + 8$ (-3, -1) **f** $y = 4x + x^{-2}$ $(\frac{1}{2}, 6)$
- 3 Evaluate $f'(4)$ when
- a** $f(x) = (x + 1)^2$ **b** $f(x) = x^{\frac{1}{2}}$ **c** $f(x) = x - 4x^{-2}$ **d** $f(x) = 5 - 6x^{\frac{3}{2}}$
- 4 The curve with equation $y = x^3 - 4x^2 + 3x$ crosses the x -axis at the points A , B and C .
- a** Find the coordinates of the points A , B and C .
- b** Find the gradient of the curve at each of the points A , B and C .
- 5 For the curve with equation $y = 2x^2 - 5x + 1$,
- a** find $\frac{dy}{dx}$,
- b** find the value of x for which $\frac{dy}{dx} = 7$.
- 6 Find the coordinates of the points on the curve with the equation $y = x^3 - 8x$ at which the gradient of the curve is 4.
- 7 A curve has the equation $y = x^3 + x^2 - 4x + 1$.
- a** Find the gradient of the curve at the point $P(-1, 5)$.
- Given that the gradient at the point Q on the curve is the same as the gradient at the point P ,
- b** find, as exact fractions, the coordinates of the point Q .
- 8 Find an equation of the tangent to each curve at the given point.
- a** $y = x^2$ (2, 4) **b** $y = x^2 + 3x + 4$ (-1, 2)
- c** $y = 2x^2 - 6x + 8$ (1, 4) **d** $y = x^3 - 4x^2 + 2$ (3, -7)
- 9 Find an equation of the tangent to each curve at the given point. Give your answers in the form $ax + by + c = 0$, where a , b and c are integers.
- a** $y = 3 - x^2$ (-3, -6) **b** $y = \frac{2}{x}$ (2, 1)
- c** $y = 2x^2 + 5x - 1$ $(\frac{1}{2}, 2)$ **d** $y = x - 3\sqrt{x}$ (4, -2)
- 10 Find an equation of the normal to each curve at the given point. Give your answers in the form $ax + by + c = 0$, where a , b and c are integers.
- a** $y = x^2 - 4$ (1, -3) **b** $y = 3x^2 + 7x + 7$ (-2, 5)
- c** $y = x^3 - 8x + 4$ (2, -4) **d** $y = x - \frac{6}{x}$ (3, 1)



- 11 Find, in the form $y = mx + c$, an equation of
- the tangent to the curve $y = 3x^2 - 5x + 2$ at the point on the curve with x -coordinate 2,
 - the normal to the curve $y = x^3 + 5x^2 - 12$ at the point on the curve with x -coordinate -3 .
- 12 A curve has the equation $y = x^3 + 3x^2 - 16x + 2$.
- Find an equation of the tangent to the curve at the point $P(2, -10)$.
The tangent to the curve at the point Q is parallel to the tangent at the point P .
 - Find the coordinates of the point Q .
- 13 A curve has the equation $y = x^2 - 3x + 4$.
- Find an equation of the normal to the curve at the point $A(2, 2)$.
The normal to the curve at A intersects the curve again at the point B .
 - Find the coordinates of the point B .
- 14 $f(x) \equiv x^3 + 4x^2 - 18$.
- Find $f'(x)$.
 - Show that the tangent to the curve $y = f(x)$ at the point on the curve with x -coordinate -3 passes through the origin.
- 15 The curve C has the equation $y = 6 + x - x^2$.
- Find the coordinates of the point P , where C crosses the positive x -axis, and the point Q , where C crosses the y -axis.
 - Find an equation of the tangent to C at P .
 - Find the coordinates of the point where the tangent to C at P meets the tangent to C at Q .
- 16 The straight line l is a tangent to the curve $y = x^2 - 5x + 3$ at the point A on the curve.
Given that l is parallel to the line $3x + y = 0$,
- find the coordinates of the point A ,
 - find the equation of the line l in the form $y = mx + c$.
- 17 The line with equation $y = 2x + k$ is a normal to the curve with equation $y = \frac{16}{x^2}$.
Find the value of the constant k .
- 18 A ball is thrown vertically downwards from the top of a cliff. The distance, s metres, of the ball from the top of the cliff after t seconds is given by $s = 3t + 5t^2$.
Find the rate at which the distance the ball has travelled is increasing when
- $t = 0.6$,
 - $s = 54$.
- 19 Water is poured into a vase such that the depth, h cm, of the water in the vase after t seconds is given by $h = kt^{\frac{1}{3}}$, where k is a constant. Given that when $t = 1$, the depth of the water in the vase is increasing at the rate of 3 cm per second,
- find the value of k ,
 - find the rate at which h is increasing when $t = 8$.

