

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--

Pearson Edexcel Level 3 GCE**Time** 2 hours**Paper
reference****8MA0/01****Mathematics****Advanced Subsidiary****PAPER 1: Pure Mathematics****You must have:**

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear.
Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 14 questions in this question paper. The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

1. Find

$$\int \left(8x^3 - \frac{3}{2\sqrt{x}} + 5 \right) dx$$

giving your answer in simplest form.

(4)

Question 1 continued

(Total for Question 1 is 4 marks)

2.

$$f(x) = 2x^3 + 5x^2 + 2x + 15$$

(a) Use the factor theorem to show that $(x + 3)$ is a factor of $f(x)$.

(2)

(b) Find the constants a , b and c such that

$$f(x) = (x + 3)(ax^2 + bx + c)$$

(2)

(c) Hence show that $f(x) = 0$ has only one real root.

(2)

(d) Write down the real root of the equation $f(x - 5) = 0$

(1)

Question 2 continued

Question 2 continued

Question 2 continued

(Total for Question 2 is 7 marks)

3. The triangle PQR is such that $\vec{PQ} = 3\mathbf{i} + 5\mathbf{j}$ and $\vec{PR} = 13\mathbf{i} - 15\mathbf{j}$

(a) Find $\vec{QR}$ (2)

(b) Hence find $|\vec{QR}|$ giving your answer as a simplified surd. (2)

The point S lies on the line segment QR so that $QS:SR = 3:2$

(c) Find $\vec{PS}$ (2)

Question 3 continued

(Total for Question 3 is 6 marks)

4.


**Figure 1**

Figure 1 shows a sketch of triangle ABC with $AB = (x + 2) \text{ cm}$, $BC = (3x + 10) \text{ cm}$, $AC = 7x \text{ cm}$, angle $BAC = 60^\circ$ and angle $ACB = \theta^\circ$

(a) (i) Show that $17x^2 - 35x - 48 = 0$ (3)

(ii) Hence find the value of x . (1)

(b) Hence find the value of θ giving your answer to one decimal place. (2)

Question 4 continued

[illegible]

Question 4 continued

Question 4 continued

(Total for Question 4 is 6 marks)

5. The mass, A kg, of algae in a small pond, is modelled by the equation

$$A = pq^t$$

where p and q are constants and t is the number of weeks after the mass of algae was first recorded.

Data recorded indicates that there is a linear relationship between t and $\log_{10} A$ given by the equation

$$\log_{10} A = 0.03t + 0.5$$

- (a) Use this relationship to find a complete equation for the model in the form

$$A = pq^t$$

giving the value of p and the value of q each to 4 significant figures.

(4)

- (b) With reference to the model, interpret

(i) the value of the constant p ,

(ii) the value of the constant q .

(2)

- (c) Find, according to the model,

(i) the mass of algae in the pond when $t = 8$, giving your answer to the nearest 0.5 kg,

(ii) the number of weeks it takes for the mass of algae in the pond to reach 4 kg.

(3)

- (d) State one reason why this may not be a realistic model in the long term.

(1)

Question 5 continued

Question 5 continued

[illegible]

Question 5 continued

(Total for Question 5 is 10 marks)

6. (a) Find the first 4 terms, in ascending powers of x , of the binomial expansion of

$$\left(3 - \frac{2x}{9}\right)^8$$

giving each term in simplest form.

(4)

$$f(x) = \left(\frac{x-1}{2x}\right)\left(3 - \frac{2x}{9}\right)^8$$

- (b) Find the coefficient of x^2 in the series expansion of $f(x)$, giving your answer as a simplified fraction.

(2)

Question 6 continued

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question 6 continued

[illegible]

Question 6 continued

(Total for Question 6 is 6 marks)

Question 7 continued

Question 7 continued

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question 7 continued

(Total for Question 7 is 7 marks)

Question 8 continued

(Total for Question 8 is 6 marks)

9. (a) Given that $p = \log_3 x$, where $x > 0$, find in simplest form in terms of p ,

(i) $\log_3\left(\frac{x}{9}\right)$

(ii) $\log_3(\sqrt{x})$

(2)

- (b) Hence, or otherwise, solve

$$2\log_3\left(\frac{x}{9}\right) + 3\log_3(\sqrt{x}) = -11$$

giving your answer as a simplified fraction.

Solutions relying on calculator technology are not acceptable.

(4)

Question 9 continued

(Total for Question 9 is 6 marks)

10.




Figure 2

In this question you must show all stages of your working.

Solutions relying on calculator technology are not acceptable.

Figure 2 shows a sketch of part of the curve C with equation

$$y = \frac{1}{3}x^2 - 2\sqrt{x} + 3 \quad x \geq 0$$

The point P lies on C and has x coordinate 4

The line l is the tangent to C at P .

(a) Show that l has equation

$$13x - 6y - 26 = 0 \quad (5)$$

The region R , shown shaded in Figure 2, is bounded by the y -axis, the curve C , the line l and the x -axis.

(b) Find the exact area of R . (5)

Question 10 continued

Question 10 continued

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question 10 continued

(Total for Question 10 is 10 marks)

Question 11 continued

Question 11 continued

Question 11 continued

(Total for Question 11 is 8 marks)

12. A company makes drinks containers out of metal.

The containers are modelled as closed cylinders with base radius r cm and height h cm and the capacity of each container is 355 cm^3

The metal used

- for the circular base and the curved side costs 0.04 pence/cm^2
- for the circular top costs 0.09 pence/cm^2

Both metals used are of negligible thickness.

(a) Show that the total cost, C pence, of the metal for one container is given by

$$C = 0.13\pi r^2 + \frac{28.4}{r} \quad (4)$$

(b) Use calculus to find the value of r for which C is a minimum, giving your answer to 3 significant figures.

(4)

(c) Using $\frac{d^2C}{dr^2}$ prove that the cost is minimised for the value of r found in part (b).

(2)

(d) Hence find the minimum value of C , giving your answer to the nearest integer.

(2)

Question 12 continued

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question 12 continued

Question 12 continued

(Total for Question 12 is 12 marks)

13. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

(a) Show that

$$\frac{1}{\cos \theta} + \tan \theta \equiv \frac{\cos \theta}{1 - \sin \theta} \quad \theta \neq (2n + 1)90^\circ \quad n \in \mathbb{Z} \quad (3)$$

Given that $\cos 2x \neq 0$

(b) solve for $0 < x < 90^\circ$

$$\frac{1}{\cos 2x} + \tan 2x = 3 \cos 2x$$

giving your answers to one decimal place.

(5)

Question 13 continued

Question 13 continued

[illegible]

Question 13 continued

(Total for Question 13 is 8 marks)

14. (i) A student states

“if x^2 is greater than 9 then x must be greater than 3”

Determine whether or not this statement is true, giving a reason for your answer.

(1)

(ii) Prove that for all positive integers n ,

$$n^3 + 3n^2 + 2n$$

is divisible by 6

(3)

Question 14 continued

Question 14 continued

(Total for Question 14 is 4 marks)

TOTAL FOR PAPER IS 100 MARKS