

Question Number	Scheme	Marks
1. (a)(i)	$X \sim B(50, 0.4)$ $P(X = 26) = 0.9686 - 0.9427$ or ${}^{50}C_{26} (0.4)^{26}(0.6)^{24}$ awrt <b><u>0.0259</u></b>	M1 A1 (2)
(ii)	$P(X \geq 26) = 1 - P(X \leq 25)$ $= 1 - 0.9427 =$ awrt <b><u>0.0573</u></b>	M1 A1 (2)
(iii)	(From tables) $k =$ <b><u>19</u></b>	B1 (1)
(b)(i)	$J \sim N(240, 144)$ $P(X \leq 222) \sim P(J < 222.5) = P\left(Z < \frac{222.5 - 240}{\sqrt{144}}\right)$ $P(Z < -1.46) = 1 - 0.9279 =$ awrt <b><u>0.0721 - 0.0724</u></b>	M1A1 M1M1 A1 (5)
(ii)	$n$ is large (oe) and $p$ is close to 0.5	B1 (1)
<b>[11 marks]</b>		
<b>Notes</b>		
(a)(i)	M1 Use of tables or ${}^{50}C_{26}(p)^{26}(1-p)^{24}$ with $0 < p < 1$ allow alternative notations for ${}^{50}C_{26}$ A1 awrt 0.0259 (correct answer scores 2 out of 2)	
(ii)	M1 writing or using $1 - P(X \leq 25)$ A1 awrt 0.0573 (calc 0.0573437....) (correct answer scores 2 out of 2)	
(iii)	B1 19 cao $k \leq 19$ or $k \geq 19$ is B0	
(b)(i)	1 <sup>st</sup> M1 For writing or using $N(240, \dots)$ (May be seen in standardisation) 1 <sup>st</sup> A1 For writing or using $N(240, 144)$ (May be seen in standardisation) 2 <sup>nd</sup> M1 use of continuity correction $222 \pm 0.5$ 3 <sup>rd</sup> M1 $\pm \left( \frac{222 \text{ or } 222.5 \text{ or } 221.5 - \text{their mean}}{\text{their sd}} \right)$ if distribution not clearly stated, then the mean and sd must be correct in the standardisation to score this mark 2 <sup>nd</sup> A1 awrt 0.0721 through to awrt 0.0724 (calc 0.0723743....) Answer in the range implies all previous marks unless clearly comes from wrong method [NB: Use of binomial distribution gives 0.0719]	
(ii)	B1 both conditions required for $n$ is large allow in words e.g. 'sample is large' allow 0.4 in place of $p$ condone ' $n > 30$ ' (or any number $> 30$ ) Ignore comments about $np$	

Question Number	Scheme	Marks
2. (a)	e.g. Population is small	B1 (1)
(b)(i)	list/register/database of <b>all</b> members (of the leisure centre)	B1
(ii)	A member (of the leisure centre)	B1 (2)
(c)	C is the statistic as it is (a quantity) based only on <u>values</u> (oe) taken from the <u>sample</u> /it contains <u>no unknown parameters/population values</u>	B1 (1)
		<b>[4 marks]</b>
	<b>Notes</b>	
(a)	B1 any correct characteristic of the population that makes a census a practical alternative to a sample (accessible, finite, well-defined)	
(b)(i)	B1 idea of list (oe) <u>and</u> idea of <b>all</b> members (e.g. list of each member of the leisure centre))	
(ii)	B1 a single member Condone members Also condone One of the members in the sample The opinion/view of one of the members is B0	
(c)	B1 choosing C (or clearly identifying C in words) only with a correct supporting reason which must include value (oe) and sample <u>or</u> no unknown parameters For values allow e.g. information, observations, calculations, function, numerical data, etc.	

Question Number	Scheme	Marks
3. (a)	$\int_2^5 \frac{1}{48}(x^2 - 8x + c) dx = 1$ $1 = \frac{1}{48} \left[ \frac{x^3}{3} - 4x^2 + cx \right]_2^5$ $1 = \frac{1}{48} \left( \left( \frac{5^3}{3} - 4(5^2) + 5c \right) - \left( \frac{2^3}{3} - 4(2^2) + 2c \right) \right) \text{ or } 48 = 39 - 84 + 3c$ $(\Rightarrow 3c = 93 \Rightarrow) c = 31^*$	M1 M1 A1cso* (3)
(b)	$P(2 < X < 3) = \frac{1}{48} \left[ \frac{x^3}{3} - 4x^2 + 31x \right]_2^3$ $\frac{1}{48} \left( \left( \frac{3^3}{3} - 4(3^2) + 31(3) \right) - \left( \frac{2^3}{3} - 4(2^2) + 31(2) \right) \right) = \frac{13}{36} \text{ (=awrt 0.361)}$	M1 A1 (2)
(c)	Less than 3 since " $\frac{13}{36}$ " > 0.25	B1 (1)
(d)	$x = 4$ leads to the minimum/lowest value of $f(x)$ / $f(x)$ is a positive quadratic	B1 (1)
(e)	Considers $x = 2$ and $x = 5$ by e.g. <ul style="list-style-type: none"> <li><math>f(2) = 0.39(58\dot{3}) [= \frac{19}{48}]</math> and <math>f(5) = 0.\dot{3} [= \frac{16}{48}]</math> (so <math>f(2) &gt; f(5)</math>)</li> <li>Sketch of <math>f(x)</math> from <math>x = 2</math> to <math>x = 5</math></li> <li><math>x = 2</math> is further than <math>x = 4</math> (then <math>x = 5</math>)</li> </ul> Mode is $x = 2$	M1 A1 (2)
<b>[9 marks]</b>		
<b>Notes</b>		
(a)	1 <sup>st</sup> M1 setting up integral and equating to 1 (condone missing dx) limits not needed 2 <sup>nd</sup> M1 attempting to integrate $f(x)$ at least one term $x^n \rightarrow x^{n+1}$ (need not be = 1) Use of integration of $f(x)$ with $F(2) = 0$ and $F(5) = 1$ can score M1M1 A1* cso including use of correct limits. There should be at least one line of working between scoring the 2 <sup>nd</sup> M1 and arriving at the given answer. Allow a verification method 1 <sup>st</sup> M1 setting up integral 2 <sup>nd</sup> M1 attempting to integrate A1cso use of correct limits to show that it integrates to 1 and concluding that $c = 31$	
(b)	M1 for use of integration of $f(x)$ $x^n \rightarrow x^{n+1}$ with correct limits 2 and 3 (ft from their (a)) A1 allow awrt 0.361 (correct answer scores 2 out of 2)	
(c)	B1 less than 3 with correct reasoning. May use their part (b), but must be consistent with 'less than 3' If the lower quartile is found awrt 2.67, allow $LQ/2.67 < 3$	
(d)	B1 correct reason why the method does not give the correct mode. Allow a sketch of $f(x)$ . Also allow, e.g. 'Kei's method did not consider the end-points'	
(e)	M1 considers end-points A1 mode is 2 cao Answer only scores M0A0. Must have some justification.	

Question Number	Scheme	Marks
4. (a)	$p$ is small	B1 (1)
(b)	Let $N$ = number of candles <b>not</b> suitable for sale $N \sim B(125, 0.02)$ $\approx C \sim \text{Po}(2.5)$ $P(C \leq 6)$ $= 0.9858$ awrt <b><u>0.986</u></b>	M1 A1 M1 A1 (4)
(c)(i)	$H_0 : p = 0.05$ $H_1 : p < 0.05$ $D \sim B(30, 0.05)$ $P(D = 0) = 0.2146$ Do not reject $H_0$ / not significant The <u>manufacturer's</u> claim <b>is not</b> supported/There <b>is not</b> enough evidence to suggest that the <u>proportion</u> (oe) of candle <u>holders</u> with minor <u>defects</u> is less than 5%/ <u>Charlie's</u> claim <b>is</b> supported	B1 M1 A1 M1 A1 (5)
(ii)	Impossible to reject $H_0$ (since $P(D = 0) > 0.05$ )	B1 (1)
(d)	$0.95^{50} [=0.0769\dots]$ .... <u>or</u> $X \sim B(50, 0.05)$ , $P(X = 0)$ (is still) $> 0.05$ (so still not possible to reject $H_0$ ) hence Ashley's change does not make the test appropriate.	M1 A1 (2)
		<b>[13 marks]</b>
	<b>Notes</b>	
(a)	B1 correct condition allow ' $p$ is close to 0' allow ' $p < 0.1$ ' or any value less than 0.1 (condone $np < 10$ or $np \leq 10$ )	
(b)	1 <sup>st</sup> M1 recognising Binomial distribution (may be implied by Po(2.5)) 1 <sup>st</sup> A1 correct distribution Po(2.5) 2 <sup>nd</sup> M1 writing or using $P(C \leq 6)$ from Poisson distribution 2 <sup>nd</sup> A1 awrt 0.986 from correct distribution used (calc : 0.9858126....) [NB : Use of binomial gives 0.98678...] Answer only 0.9858 or better scores 4 out of 4, but answer of 0.986 must see Po(2.5) to award full marks.	
(c)(i)	B1 correct hypotheses in terms of $p$ or $\pi$ 1 <sup>st</sup> M1 writing or using $B(30, 0.05)$ (may be implied by 1 <sup>st</sup> A1) 1 <sup>st</sup> A1 awrt 0.215 2 <sup>nd</sup> M1 a correct ft statement consistent with their $p$ -value and 0.05 No context needed but do not allow contradicting non contextual comments. 2 <sup>nd</sup> A1 correct conclusion in context which must be <b>not rejecting</b> $H_0$ . Must use underlined words (oe) No hypotheses then A0 Condone e.g. '5% of candle holders have minor defects'	
(ii)	B1 correct reasoning which implies there is no critical region/ $H_0$ cannot be rejected Sample size is too small on its own is B0.	
(d)	M1 for $0.95^{50}$ <u>or</u> for $X \sim B(50, 0.05)$ and $P(X = 0) > 0.05$ A1 test is (still) not appropriate with M1 scored	

Question Number	Scheme	Marks
5. (a)	$F(3) = 0 \rightarrow \frac{1}{16}(3^2 - 6(3) + a) = 0$ $a = 9$ $F(10) = 1 \rightarrow \frac{1}{12}(100(10) - (5)10^2 + c) = 1$ $c = -488$	M1 A1 M1 A1 (4)
(b)	$\frac{1}{16}(5^2 - 6(5) + "9") = \frac{1}{12}(5 + b) \quad \left  \quad \frac{1}{12}(9 + b) = \frac{1}{12}(100(9) - 5(9^2) + "-488") \right.$ $b = -2$	M1 A1 (2)
(c)	$P(6 < Y \leq 9) = F(9) - F(6)$ $= \frac{1}{12}(9 + "-2") - \frac{1}{12}(6 + "-2")$ $= \frac{1}{4}$	M1 M1 A1 (3)
(d)	$f(y) = \frac{1}{12}$	B1 (1)
(e)	$E(6Y - 5) = [26.5 + ] \int_5^9 (6y - 5) \frac{1}{12} dy$ $= [26.5 + ] \frac{1}{12} [(3y^2 - 5y)]_5^9$ $= 26.5 + \frac{1}{12} [(3(9^2) - 5(9)) - (3(5^2) - 5(5))]$ $= \frac{233}{6}$	M1 dM1 dM1 A1 (4)
<b>[Total 14]</b>		
<b>Notes</b>		
(a)	1 <sup>st</sup> M1 writing or use of $F(3) = 0$ 1 <sup>st</sup> A1 $a = 9$ cao 2 <sup>nd</sup> M1 writing or use of $F(10) = 1$ 2 <sup>nd</sup> A1 $c = -488$ cao	
(b)	M1 use of $F(5) = F(5) [= \frac{1}{4}]$ or $F(9) = F(9) [= \frac{7}{12}]$ ft their values from (a) A1 $b = -2$ cao	
(c)	1 <sup>st</sup> M1 writing or using $F(9) - F(6)$ (may be implied by 2 <sup>nd</sup> M1) 2 <sup>nd</sup> M1 substituting 9 and 6 into $F(x)$ with their value of $b$ allow $\frac{1}{12}(100(9) + 5(9^2) + "-488") - \frac{1}{12}(6 + "-2")$ with their value of $b$ and their value of $c$ A1 $\frac{1}{4}$ oe	
(d)	B1 $\frac{1}{12}$	
(e)	1 <sup>st</sup> M1 use of $\int_5^9 (6y - 5) \frac{1}{12} dy$ (ignore limits) 2 <sup>nd</sup> M1 (dep on 1 <sup>st</sup> M1) attempt to integrate $(6y - 5) \frac{1}{12}$ with at least one $y^n \rightarrow y^{n+1}$ 3 <sup>rd</sup> M1 (dep on 1 <sup>st</sup> M1) $26.5 + \int_5^9 (6y - 5) \frac{1}{12} dy$ A1 awrt 38.8	
SC:	Answer only or correct answer not using given information scores M0M1M1A1	

Question Number	Scheme	Marks
<b>6. (a)</b>	$P(17 < W < k) = P(W < k) - P(W < 17) = \frac{53}{60} - \left(1 - \frac{1}{5}\right) = \frac{1}{12}$	M1 A1 (2)
<b>(b)(i)</b>	$\frac{(b-a)^2}{12} = 75, \quad \frac{b-17}{b-a} = \frac{1}{5} \quad \text{or} \quad \frac{17-a}{b-a} = \frac{4}{5}$ $\frac{(b-a)^2}{12} = 75 \rightarrow (b-a) = 30 \quad \frac{b-17}{30} = \frac{1}{5}$ $b = 23 \text{ and } a = -7$	B1, B1 M1 A1 (4)
<b>(ii)</b>	$P(W < k) = \frac{k - (" - 7")}{"23" - (" - 7")} = \frac{53}{60} \quad \text{or} \quad P(17 < W < k) = \frac{k-17}{30} = \frac{1}{12} \quad \text{or} \quad P(W > k) = \frac{"23" - k}{"23" - (" - 7")} = \frac{7}{60}$ $k = 19.5$	M1 A1 (2)
<b>(c)</b>	$P(-5 < W < 5) = \frac{5 - (-5)}{"23" - (" - 7")} = \frac{1}{3}$	M1A1ft (2)
<b>(d)</b>	$E(W^2) = \text{Var}(W) + E(W)^2 = 75 + \left(\frac{"23" + " - 7"}{2}\right)^2 = 139$	M1 A1 (2)
<b>[Total 12]</b>		
<b>Notes</b>		
<b>(a)</b>	M1 for writing or using $P(W < k) - P(W < 17)$ allow $<$ or $\leq$ Allow equivalent expressions e.g. $P(W > 17) - P(W > k) = \frac{1}{5} - \left(1 - \frac{53}{60}\right)$ A1 oe condone awrt 0.0833 condone $\frac{1}{12}$ coming from $\frac{13}{12} - 1$ or $\left  -\frac{1}{12} \right $	
<b>(b) (i)</b>	1 <sup>st</sup> B1 correct equation for variance 2 <sup>nd</sup> B1 either correct probability equation Allow e.g. $k$ in place of $(b - a)$ 1 <sup>st</sup> M1 eliminating $(b - a)$ which must appear in both equations. A1 both $b = 23$ and $a = -7$ correct answers imply all 4 marks	
<b>(ii)</b>	M1 probability expression using uniform distribution ft their values A1 $k = 19.5$ oe cao	
<b>(c)</b>	M1 for $10/(\text{their } b - \text{their } a)$ A1ft $\frac{1}{3}$ oe condone awrt 0.333 (Allow ft $\frac{10}{\text{their}(b-a)}$ as exact fraction or evaluated to 3sf or better provided $a < -5$ and $b > 5$ )	
<b>(d)</b>	M1 use of $E(W^2) = \text{Var}(W) + (E(W))^2$ with values substituted for $\text{Var}(W)$ and $E(W)$ ft their values of $a$ and $b$ allow any rearrangement. Must have a correct (ft) expression or value for $E(W)$ Also allow $\int_{"-7"}^{"23"} \frac{1}{"23" - " - 7"} w^2 dw$ A1 139 cao	

Question Number	Scheme	Marks
7. (a)	$R \sim \text{Po}(8)$ $P(4 \leq R \leq 8) = P(R \leq 8) - P(R \leq 3) = 0.5925 - 0.0424$ $= 0.5501 = \text{awrt } \underline{0.550}$	B1 M1 A1 (3)
(b)	$H \sim \text{Po}(4)$ $P(H \leq 2) = 0.2381$ $Y \sim B(5, "0.2381")$ $P(Y = 2) = {}^5C_2 ("0.2381")^2 (1 - "0.2381")^3$ $= 0.25073\dots = \text{awrt } \underline{0.251}$	B1 B1 M1 M1 A1 (5)
(c)	$W = \text{number sold in first fifteen minutes}$ $X = \text{number sold in last forty five minutes}$ $P(W > X \mid R = 4) = \frac{P(W = 4)P(X = 0) + P(W = 3)P(X = 1)}{P(R = 4)}$ $= \frac{\frac{e^{-2} 2^4}{4!} \frac{e^{-6} 6^0}{0!} + \frac{e^{-2} 2^3}{3!} \frac{e^{-6} 6^1}{1!}}{\frac{e^{-8} 8^4}{4!}}$	$F = \text{number of muffins sold in first 15 minutes}$ $F \sim B(4, 0.25)$ $P(F > 2) =$ $P(F = 3) + P(F = 4)$ $= {}^4C_3 (0.25)^3 (0.75) + 0.25^4$ $= \frac{13}{256} (\text{awrt } 0.0508 \text{ or awrt } 0.0509)$ M1 M1 M1 A1 (4)
[Total 12]		
Notes		
(a)	B1 writing or using Po(8) (may be implied by one correct probability from 0.5925, 0.0424 0.4530 or 0.0996) M1 writing or using $P(R \leq 8) - P(R \leq 3)$ A1 awrt 0.550 (calc: 0.55016....) correct answer scores 3 out of 3	
(b)	1 <sup>st</sup> B1 writing or using Po(4) 2 <sup>nd</sup> B1 awrt 0.238 1 <sup>st</sup> M1 choosing binomial distribution with $n = 5$ and their $p$ 2 <sup>nd</sup> M1 ${}^5C_2 p^2(1-p)^3$ with $0 < p < 1$ A1 awrt 0.251	
(c)	1 <sup>st</sup> M1 attempt at either correct product $P(W = 4)P(X = 0)$ or $P(W = 3)P(X = 1)$ from $W \sim \text{Po}(2)$ and $X \sim \text{Po}(6)$ implied by awrt 0.0902×awrt 0.0025 or awrt 0.180×awrt 0.0149 or awrt 0.0029 2 <sup>nd</sup> M1 conditional probability with $P(R = 4)$ from $R \sim \text{Po}(8)$ on denominator implied by awrt 0.0573 seen in the denominator of a probability expression 3 <sup>rd</sup> M1 complete expression for the required probability implied (awrt 0.0902×awrt 0.0025+awrt 0.180×awrt 0.0149)/awrt 0.0573 for 3 <sup>rd</sup> M1 A1 allow awrt 0.0508 or awrt 0.0509 from use of tables	
ALT	1 <sup>st</sup> M1 identifying B(4, 0.25) 2 <sup>nd</sup> M1 $P(F = 3) + P(F = 4)$ from B(4, 0.25) 3 <sup>rd</sup> M1 $4p^3q + p^4$ from B(4, 0.25)	