Solomon Practice Paper

Pure Mathematics 2J

Time allowed: 90 minutes

Centre: www.CasperYC.club

Name:

Teacher:

Question	Points	Score
1	4	
2	8	
3	8	
4	10	
5	11	
6	11	
7	11	
8	12	
Total:	75	

How I can achieve better:

•

•

•





[4]

1.	Given that $y = 3e^x + 2\ln(x)$, find $\frac{d^2y}{dx^2}$.
	passag

2.	(a) By letting $p = \log_a(x)$ and $q = \log_a(y)$, or otherwise, prove that	[4]
	$\log_a(xy) \equiv \log_a(x) + \log_a(y).$	
	(b) Find integers A and B such that	[4]
	$\ln(48) + \ln(108) = A\ln(2) + B\ln(3).$	
	,	Total: 8

3.	(a) Express $\left(x^{\frac{1}{2}} - 2x^{-\frac{3}{2}}\right)^2$ in the form $px + qx^{-1} + rx^{-3}$. (b) Show that	[3] [5]
	$\int_{2}^{4} \left(x^{\frac{1}{2}} - 2x^{-\frac{3}{2}} \right)^{2} dx = \frac{51}{8} - 4\ln(2).$	
		Total: 8
		10tai: 8

4.	(a) Find the values of θ in the interval $0 \le \theta \le 2\pi$, for which	[5
	$2\tan^2(\theta) + \sec^2(\theta) = 2,$	
	giving your answers in terms of π .	
	(b) Find the values of x in the interval $0 \le x \le 180^{\circ}$, for which	[5
		-
	$\sin(3x) = \sin(2x).$	
	r	Fotal: 10
		rotar. r

Last updated: May 5, 2023

5.	Given that $a > 0$,	
	(a) sketch on the same set of coordinate axes the graphs of $y = \frac{1}{2}(x+a)$ and $y = 2x-a $, labelling the coordinates of any points where each graph meets the coordinate axes,	[5]
	(b) find, in terms of a, the coordinates of any points where the two graphs intersect.	[6]
		Total: 11

Last updated: May 5, 2023



6.	(a) Expand $(4+2x)^5$ as a series in ascending powers of x, simplifying each coefficient.	[4]
	Hence, find	
	(b) the coefficient of y^4 in the expansion of $(4 + \frac{1}{5}y)^5$ as an exact fraction,	[3]
	(c) the coefficient of z^6 in the expansion of $(2+\sqrt{2}z)^5(2-\sqrt{2}z)^5$.	[4]
		Total: 11
		10001. 11
		•
		,
		,
		•
		•
		•
		,
		,
		•
		,
		,
		•
		•
		•
		•
		•

[2]

[2]

Total: 11

7.

$$f(x) \equiv x^4 - 5x + 3.$$

- (a) Show that one root of the equation f(x) = 0 lies in the interval (0.6, 0.7). [2]
- (b) Using the iteration formula [3]

$$x_{n+1} = 0.2 \left(x_n^4 + 3 \right),\,$$

with a starting value of $x_1 = 0.65$, find this root correct to 3 significant figures.

(c) Show that the equation f(x) = 0 can be rewritten as

$$x = \pm \sqrt{\frac{ax+b}{x^2}}$$

where a and b are integers to be found.

(d) Hence, use the iteration formula

$$x_{n+1} = \pm \sqrt{\frac{a_n x + b}{x_n^2}},$$

together with your values of a and b and with $x_1 = 1.5$ to find x_2, x_3 and x_4 correct to 6 significant figures.


(e) Considering only your values of x_2, x_3 and x_4 , explain why it is reasonable to give a second root of the equation as 1.43 correct to 3 significant figures.

• • • •	 • •	 	 	 	 	 	 	 	 	 • •	 	 	 	 	 	 	
	 • •	 • •	 	 	 	 	 ٠.	 	 ٠.	 • •	 	 	 • •	 	 	 	
	 ٠.	 	 	 	 	 	 	 	 	 	 	 	 	 	 	 	



		 	• • • • •
•••••		 	
		 	• • • • •
•••••		 	• • • • •
			• • • • •
	• • • • • • • • • • • • • • • • • • • •	 	• • • • •
			• • • • •
			• • • • •
			• • • • •
		 	• • • • •
• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	 	

8. Figure shows the straight line l and the curve y = f(x).



The line and curve intersect at the points $P(1, \ln(2))$ and $Q(3, \ln(8))$.

(a) Find in its simplest form the equation of the line l.

[4]

Given that $f(x) \equiv \ln(ax + b)$,

(b) find the values of a and b,

[5]

[3]

(c) hence, find an expression for $f^{-1}(x)$.

Total: 12

• • • • • • • • • • • • • • • • • • • •	 	

 	• • • • • • • • • • • • • • • • • • • •	

