


THERE ARE 14 QUESTIONS IN THIS PAPER.

Q1 Q2 Q3 Q4 Q5 Q6 Q7 Q8 Q9 Q10 Q11 Q12 Q13 Q14

Sixth Term Examination Paper

04-S3



Compiled by: Dr Yu 郁博士

www.CasperYC.club

Last updated: May 8, 2025



SUGGESTIONS TO DRYUFROMSHANGHAI@QQ.COM

Section A: Pure Mathematics

1 (i) Show that

$$\int_0^a \frac{\sinh x}{2 \cosh^2 x - 1} dx = \frac{1}{2\sqrt{2}} \ln \left(\frac{\sqrt{2} \cosh a - 1}{\sqrt{2} \cosh a + 1} \right) + \frac{1}{2\sqrt{2}} \ln \left(\frac{\sqrt{2} + 1}{\sqrt{2} - 1} \right)$$

and find

$$\int_0^a \frac{\cosh x}{1 + 2 \sinh^2 x} dx.$$

(ii) Hence show that

$$\int_0^\infty \frac{\cosh x - \sinh x}{1 + 2 \sinh^2 x} dx = \frac{\pi}{2\sqrt{2}} - \frac{1}{2\sqrt{2}} \ln \left(\frac{\sqrt{2} + 1}{\sqrt{2} - 1} \right).$$

(iii) By substituting $u = e^x$ in this result, or otherwise, find

$$\int_1^\infty \frac{1}{1 + u^4} du.$$

2 The equation of a curve is $y = f(x)$ where

$$f(x) = x - 4 - \frac{16(2x + 1)^2}{x^2(x - 4)}.$$

(i) Write down the equations of the vertical and oblique asymptotes to the curve and show that the oblique asymptote is a tangent to the curve.

(ii) Show that the equation $f(x) = 0$ has a double root.

(iii) Sketch the curve.

- 3 (i) Given that $f''(x) > 0$ when $a \leq x \leq b$, explain with the aid of a sketch why

$$(b-a)f\left(\frac{a+b}{2}\right) < \int_a^b f(x) \, dx < (b-a) \frac{f(a)+f(b)}{2}.$$

- (ii) By choosing suitable a, b and $f(x)$, show that

$$\frac{4}{(2n-1)^2} < \frac{1}{n-1} - \frac{1}{n} < \frac{1}{2} \left(\frac{1}{n^2} + \frac{1}{(n-1)^2} \right),$$

where n is an integer greater than 1.

- (iii) Deduce that

$$4 \left(\frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots \right) < 1 < \frac{1}{2} + \left(\frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots \right).$$

- (iv) Show that

$$\frac{1}{2} \left(\frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \frac{1}{6^2} + \dots \right) < \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots$$

and hence show that

$$\frac{3}{2} < \sum_{n=1}^{\infty} \frac{1}{n^2} < \frac{7}{4}.$$

- 4 The triangle OAB is isosceles, with $OA = OB$ and angle $AOB = 2\alpha$ where $0 < \alpha < \frac{\pi}{2}$. The semi-circle C_0 has its centre at the midpoint of the base AB of the triangle, and the sides OA and OB of the triangle are both tangent to the semi-circle. $C_1, C_2, C_3, \dots$ are circles such that C_n is tangent to C_{n-1} and to sides OA and OB of the triangle.

- (i) Let r_n be the radius of C_n . Show that

$$\frac{r_{n+1}}{r_n} = \frac{1 - \sin \alpha}{1 + \sin \alpha}.$$

- (ii) Let S be the total area of the semi-circle C_0 and the circles $C_1, C_2, C_3, \dots$. Show that

$$S = \frac{1 + \sin^2 \alpha}{4 \sin \alpha} \pi r_0^2.$$

- (iii) Show that there are values of α for which S is more than four fifths of the area of triangle OAB .

- 5 Show that if $\cos(x - \alpha) = \cos \beta$ then either $\tan x = \tan(\alpha + \beta)$ or $\tan x = \tan(\alpha - \beta)$.

By choosing suitable values of x, α and β , give an example to show that if $\tan x = \tan(\alpha + \beta)$, then $\cos(x - \alpha)$ need not equal $\cos \beta$.

Let ω be the acute angle such that $\tan \omega = \frac{4}{3}$.

- (i) For $0 \leq x \leq 2\pi$, solve the equation

$$\cos x - 7 \sin x = 5$$

giving both solutions in terms of ω .

- (ii) For $0 \leq x \leq 2\pi$, solve the equation

$$2 \cos x + 11 \sin x = 10$$

showing that one solution is twice the other and giving both in terms of ω .

- 6 Given a sequence $w_0, w_1, w_2, \dots$, the sequence $F_1, F_2, \dots$ is defined by

$$F_n = w_n^2 + w_{n-1}^2 - 4w_n w_{n-1}.$$

Show that $F_n - F_{n-1} = (w_n - w_{n-2})(w_n + w_{n-2} - 4w_{n-1})$ for $n \geq 2$.

- (i) The sequence $u_0, u_1, u_2, \dots$ has $u_0 = 1$, and $u_1 = 2$ and satisfies

$$u_n = 4u_{n-1} - u_{n-2} \quad (n \geq 2).$$

Prove that $u_n^2 + u_{n-1}^2 = 4u_n u_{n-1} - 3$ for $n \geq 1$.

- (ii) A sequence $v_0, v_1, v_2, \dots$ has $v_0 = 1$ and satisfies

$$v_n^2 + v_{n-1}^2 = 4v_n v_{n-1} - 3 \quad (n \geq 1). \quad (*)$$

(a) Find v_1 and prove that, for each $n \geq 2$, either $v_n = 4v_{n-1} - v_{n-2}$ or $v_n = v_{n-2}$.

(b) Show that the sequence, with period 2, defined by

$$v_n = \begin{cases} 1 & \text{for } n \text{ even} \\ 2 & \text{for } n \text{ odd} \end{cases}$$

satisfies (*).

(c) Find a sequence v_n with period 4 which has $v_0 = 1$, and satisfies (*).

7 For $n = 1, 2, 3, \dots$, let

$$I_n = \int_0^1 \frac{t^{n-1}}{(t+1)^n} dt.$$

(i) By considering the greatest value taken by $ds \frac{t}{t+1}$ for $0 \leq t \leq 1$ show that $I_{n+1} < \frac{1}{2} I_n$.

(ii) Show also that $ds I_{n+1} = -\frac{1}{n 2^n} + I_n$.

(iii) Deduce that $ds I_n < \frac{1}{n 2^{n-1}}$.

(iv) Prove that

$$\ln 2 = \sum_{r=1}^n \frac{1}{r 2^r} + I_{n+1}$$

and hence show that $\frac{2}{3} < \ln 2 < \frac{17}{24}$.

8 (i) Show that if

$$\frac{dy}{dx} = f(x)y + \frac{g(x)}{y}$$

then the substitution $u = y^2$ gives a linear differential equation for $u(x)$.

(ii) Hence or otherwise solve the differential equation

$$\frac{dy}{dx} = \frac{y}{x} - \frac{1}{y}.$$

(iii) Determine the solution curves of this equation which pass through $(1, 1)$, $(2, 2)$ and $(4, 4)$ and sketch graphs of all three curves on the same axes.

Section B: Mechanics

- 9** A circular hoop of radius a is free to rotate about a fixed horizontal axis passing through a point P on its circumference. The plane of the hoop is perpendicular to this axis. The hoop hangs in equilibrium with its centre, O , vertically below P . The point A on the hoop is vertically below O , so that POA is a diameter of the hoop.

A mouse M runs at constant speed u round the rough inner surface of the lower part of the hoop.

- (i) Show that the mouse can choose its speed so that the hoop remains in equilibrium with diameter POA vertical.
- (ii) Describe what happens to the hoop when the mouse passes the point at which angle $AOM = 2 \arctan \mu$, where μ is the coefficient of friction between mouse and hoop.

- 10** A particle P of mass m is attached to points A and B , where A is a distance $9a$ vertically above B , by elastic strings, each of which has modulus of elasticity $6mg$. The string AP has natural length $6a$ and the string BP has natural length $2a$. Let x be the distance AP .

- (i) The system is released from rest with P on the vertical line AB and $x = 6a$. Show that the acceleration $\ddot{x}$ of P is $\frac{4g}{a}(7a - x)$ for $6a < x < 7a$ and $\frac{g}{a}(7a - x)$ for $7a < x < 9a$.

- (ii) Find the time taken for the particle to reach B .

- 11** Particles P , of mass 2, and Q , of mass 1, move along a line. Their distances from a fixed point are x_1 and x_2 , respectively where $x_2 > x_1$. Each particle is subject to a repulsive force from the other of magnitude $\frac{2}{z^3}$, where $z = x_2 - x_1$.

- (i) Initially, $x_1 = 0, x_2 = 1$, Q is at rest and P moves towards Q with speed 1. Show that z obeys the equation $\frac{d^2z}{dt^2} = \frac{3}{z^3}$.

- (ii) By first writing $\frac{d^2z}{dt^2} = v \frac{dv}{dz}$, where $v = \frac{dz}{dt}$, show that $z = \sqrt{4t^2 - 2t + 1}$.

- (iii) By considering the equation satisfied by $2x_1 + x_2$, find x_1 and x_2 in terms of t .

Section C: Probability and Statistics

- 12** A team of m players, numbered from 1 to m , puts on a set of m shirts, similarly numbered from 1 to m . The players change in a hurry, so that the shirts are assigned to them randomly, one to each player.
- (i) Let C_i be the random variable that takes the value 1 if player i is wearing shirt i , and 0 otherwise. Show that $E(C_1) = \frac{1}{m}$ and find $\text{Var}(C_1)$ and $\text{Cov}(C_1, C_2)$.
 - (ii) Let $N = C_1 + C_2 + \cdots + C_m$ be the random variable whose value is the number of players who are wearing the correct shirt. Show that $E(N) = \text{Var}(N) = 1$.
 - (iii) Explain why a Normal approximation to N is not likely to be appropriate for any m , but that a Poisson approximation might be reasonable.
 - (iv) In the case $m = 4$, find, by listing equally likely possibilities or otherwise, the probability that no player is wearing the correct shirt and verify that an appropriate Poisson approximation to N gives this probability with a relative error of about 2%. [Use $e \approx 2\frac{72}{100}$.]
- 13** A men's endurance competition has an unlimited number of rounds. In each round, a competitor has, independently, a probability p of making it through the round; otherwise, he fails the round. Once a competitor fails a round, he drops out of the competition; before he drops out, he takes part in every round. The grand prize is awarded to any competitor who makes it through a round which all the other remaining competitors fail; if all the remaining competitors fail at the same round the grand prize is not awarded. If the competition begins with three competitors, find the probability that:
- (i) all three drop out in the same round;
 - (ii) two of them drop out in round r (with $r \geq 2$) and the third in an earlier round;
 - (iii) the grand prize is awarded.

14 In this question, $\Phi(z)$ is the cumulative distribution function of a standard normal random variable.

A random variable is known to have a Normal distribution with mean μ and standard deviation either σ_0 or σ_1 , where $\sigma_0 < \sigma_1$. The mean, $\bar{X}$, of a random sample of n values of X is to be used to test the hypothesis $H_0 : \sigma = \sigma_0$ against the alternative $H_1 : \sigma = \sigma_1$.

- (i) Explain carefully why it is appropriate to use a two sided test of the form: accept H_0 if $\mu - c < \bar{X} < \mu + c$, otherwise accept H_1 .
- (ii) Given that the probability of accepting H_1 when H_0 is true is α , determine c in terms of n, σ_0 and z_α , where z_α is defined by $\text{ds}\Phi(z_\alpha) = 1 - \frac{1}{2}\alpha$.
- (iii) The probability of accepting H_0 when H_1 is true is denoted by β . Show that β is independent of n .
- (iv) Given that $\Phi(1.960) \approx 0.975$ and that $\Phi(0.063) \approx 0.525$, determine, approximately, the minimum value of $\text{ds}\frac{\sigma_1}{\sigma_0}$ if α and β are both to be less than 0.05.