


THERE ARE 14 QUESTIONS IN THIS PAPER.

Q1 Q2 Q3 Q4 Q5 Q6 Q7 Q8 Q9 Q10 Q11 Q12 Q13 Q14

Sixth Term Examination Paper


97-S2



Compiled by: Dr Yu 郁博士

www.CasperYC.club

Last updated: May 8, 2025



SUGGESTIONS TO DRYUFROMSHANGHAI@QQ.COM

Section A: Pure Mathematics

- 1 Find the sum of those numbers between 1000 and 6000 every one of whose digits is one of the numbers 0, 2, 5 or 7, giving your answer as a product of primes.

- 2 Suppose that

$$3 = \frac{2}{x_1} = x_1 + \frac{2}{x_2} = x_2 + \frac{2}{x_3} = x_3 + \frac{2}{x_4} = \dots$$

Guess an expression, in terms of n , for x_n . Then, by induction or otherwise, prove the correctness of your guess.

- 3 (i) Find constants a , b , c and d such that

$$\frac{ax + b}{x^2 + 2x + 2} + \frac{cx + d}{x^2 - 2x + 2} = \frac{1}{x^4 + 4}.$$

- (ii) Show that

$$\int_0^1 \frac{dx}{x^4 + 4} = \frac{1}{16} \ln 5 + \frac{1}{8} \tan^{-1} 2.$$

- 4 Show that, when the polynomial $p(x)$ is divided by $(x - a)$, where a is a real number, the remainder is $p(a)$.

- (i) When the polynomial $p(x)$ is divided by $x - 1$, $x - 2$, $x - 3$ the remainders are 3, 1, 5 respectively. Given that

$$p(x) = (x - 1)(x - 2)(x - 3)q(x) + r(x),$$

where $q(x)$ and $r(x)$ are polynomials with $r(x)$ having degree less than three, find $r(x)$.

- (ii) Find a polynomial $P(x)$ of degree $n + 1$, where n is a given positive integer, such that for each integer a satisfying $0 \leq a \leq n$, the remainder when $P_n(x)$ is divided by $x - a$ is a .

- 5 The complex numbers $w = u + iv$ and $z = x + iy$ are related by the equation

$$z = (\cos v + i \sin v)e^u.$$

- (i) Find all w which correspond to $z = ie$.
- (ii) Find the loci in the x - y plane corresponding to the lines $u = \text{constant}$ in the u - v plane. Find also the loci corresponding to the lines $v = \text{constant}$. Illustrate your answers with clearly labelled sketches.
- (iii) Identify two subsets W_1 and W_2 of the u - v plane each of which is in one-to-one correspondence with the first quadrant $\{(x, y) : x > 0, y > 0\}$ of the x - y plane. Identify also two subsets W_3 and W_4 each of which is in one-to-one correspondence with the set $\{z : 0 < |z| < 1\}$.

[**NB** 'one-to-one' means here that to each value of w there is only one corresponding value of z , and vice-versa.]

- 6 (i) Show that, if $\tan^2 \phi = 2 \tan \phi + 1$, then $\tan 2\phi = -1$.

- (ii) Find all solutions of the equation

$$\tan \theta = 2 + \tan 3\theta$$

which satisfy $0 < \theta < 2\pi$, expressing your answers as rational multiples of π .

- (iii) Find all solutions of the equation the equation

$$\cot \theta = 2 + \cot 3\theta$$

which satisfy

$$-\frac{3\pi}{2} < \theta < \frac{\pi}{2}.$$

- 7 Let

$$y^2 = x^2(a^2 - x^2),$$

where a is a real constant. Find, in terms of a , the maximum and minimum values of y .

Sketch carefully on the same axes the graphs of y in the cases $a = 1$ and $a = 2$.

8 (i) If $f(t) \geq g(t)$ for $a \leq t \leq b$, explain very briefly why $\int_a^b f(t)dt \geq \int_a^b g(t)dt$.

(ii) Prove that if $p > q > 0$ and $x \geq 1$ then

$$\frac{x^p - 1}{p} \geq \frac{x^q - 1}{q}.$$

(iii) Show that this inequality also holds when $p > q > 0$ and $0 \leq x \leq 1$.

(iv) Prove that, if $p > q > 0$ and $x \geq 0$, then

$$\frac{1}{p} \left(\frac{x^p}{p+1} - 1 \right) \geq \frac{1}{q} \left(\frac{x^q}{q+1} - 1 \right).$$

Section B: Mechanics

- 9 A uniform solid sphere of diameter d and mass m is drawn slowly and without slipping from horizontal ground onto a step of height $d/4$ by a horizontal force which is always applied to the highest point of the sphere and is always perpendicular to the vertical plane which forms the face of the step.

- (i) Find the maximum horizontal force throughout the movement,
- (ii) and prove that the coefficient of friction between the sphere and the edge of the step must exceed $1/\sqrt{3}$.

- 10 *In this question the effect of gravity is to be neglected.*

A small body of mass M is moving with velocity v along the axis of a long, smooth, fixed, circular cylinder of radius L . An internal explosion splits the body into two spherical fragments, with masses qM and $(1 - q)M$, where $q \leq \frac{1}{2}$.

- (i) After bouncing perfectly elastically off the cylinder (one bounce each) the fragments collide and coalesce at a point $\frac{1}{2}L$ from the axis. Show that $q = \frac{3}{8}$.
- (ii) The collision occurs at a time $5L/v$ after the explosion. Find the energy imparted to the fragments by the explosion, and find the velocity after coalescence.

- 11 A tennis player serves from height H above horizontal ground, hitting the ball downwards with speed v at an angle α below the horizontal. The ball just clears the net of height h at horizontal distance a from the server and hits the ground a further horizontal distance b beyond the net.

- (i) Show that

$$v^2 = \frac{g(a+b)^2(1 + \tan^2 \alpha)}{2[H - (a+b)\tan \alpha]}$$

and

$$\tan \alpha = \frac{2a+b}{a(a+b)}H - \frac{a+b}{ab}h.$$

- (ii) By considering the signs of v^2 and $\tan \alpha$, find upper and lower bounds on H for such a serve to be possible.

Section C: Probability and Statistics

- 12** The game of Cambridge Whispers starts with the first participant Albert flipping an unbiased coin and whispering to his neighbour Bertha whether it fell 'heads' or 'tails'. Bertha then whispers this information to her neighbour, and so on. The game ends when the final player Zebedee whispers to Albert and the game is won, by all players, if what Albert hears is correct. The acoustics are such that the listeners have, independently at each stage, only a probability of $2/3$ of hearing correctly what is said.

- (i) Find the probability that the game is won when there are just three players.
- (ii) By considering the binomial expansion of $(a + b)^n + (a - b)^n$, or otherwise, find a concise expression for the probability P that the game is won when it is played by n players each having a probability p of hearing correctly.
- (iii) To avoid the trauma of a lost game, the rules are now modified to require Albert to whisper to Bertha what he hears from Zebedee, and so keep the game going, if what he hears from Zebedee is not correct. Find the expected total number of times that Albert whispers to Bertha before the modified game ends.

[You may use without proof the fact that $\sum_{k=1}^{\infty} kx^{k-1} = (1 - x)^{-2}$ for $|x| < 1$.]

- 13** A needle of length two cm is dropped at random onto a large piece of paper ruled with parallel lines two cm apart.

- (i) By considering the angle which the needle makes with the lines, find the probability that the needle crosses the nearest line given that its centre is x cm from it, where $0 < x < 1$.
- (ii) Given that the centre of the needle is x cm from the nearest line and that the needle crosses that line, find the cumulative distribution function for the length of the shorter segment of the needle cut off by the line.
- (iii) Find the probability that the needle misses all the lines.

- 14** Traffic enters a tunnel which is 9600 metres long, and in which overtaking is impossible. The number of vehicles which enter in any given time is governed by the Poisson distribution with mean 6 cars per minute. All vehicles travel at a constant speed until forced to slow down on catching up with a slower vehicle ahead. I enter the tunnel travelling at 30 m s^{-1} and all the other traffic is travelling at 32 m s^{-1} .

- (i) What is the expected number of vehicles in the queue behind me when I leave the tunnel?
- (ii) Assuming again that I travel at 30 m s^{-1} , but that all the other vehicles are independently equally likely to be travelling at 30 m s^{-1} or 32 m s^{-1} , find the probability that exactly two vehicles enter the tunnel within 20 seconds of my doing so and catch me up before I leave it.
- (iii) Find also the probability that there are exactly two vehicles queuing behind me when I leave the tunnel.

[Ignore the lengths of the vehicles.]